



## **Pre & Post Care for Hylenex & DePuffing Treatment**

### **Before Treatment:**

- ❖ Bring photos of the area of concern on a “good day” and “bad day”
- ❖ If you have open lesions, active cold sores, facial dermatitis or facial irruption you must wait until everything is healed

### **Considerations/Contraindications**

- ❖ Caution using hylenex if you have an allergy to insects/bee stings! Inform your provider and consider a test spot.

### **Post-Procedure**

- ❖ Some mild redness and swelling is normal for the first few days. Occasionally, the skin may look slightly dry post treatment. This normalizes in days-weeks.
- ❖ Patients should avoid strenuous exercise for 72 hours.
- ❖ We recommend waiting 2 weeks after dissolving to re-inject, if appropriate.
- ❖ The effect of the Hylenex/dissolving will begin to show between 1-36 hours from the time of treatment.

### **Hylenex**

#### *Injected dissolver only*

- ❖ Advise the provider if you have an allergy or hypersensitivity to bee stings, this is a contraindication.
- ❖ Some areas such as lips will swell more before the dissolving agent breaks down the product.
- ❖ The dissolving agent will sting slightly during injection, but reduces unwanted filler very quickly.
- ❖ Massage the area the day of injection to work it in to the area unless advised otherwise
- ❖ Keep the skin very hydrated with a gentle moisturizer and constant use of SPF post treatment.

### **DePuffing Treatment**

#### *Dissolver plus RF Microneedling and lymphatic massage*

- ❖ Advise the provider if you have any electrical device implanted in the body, this is a contraindication.
- ❖ The treatment will have 3 parts, ending with the RF microneedling
- ❖ Use gentle skincare, moisturizer and SPF post treatment. Redness and swelling is normal. It may look worse before looking better.
- ❖ Do not pick any small scabs that may occur from the RF microneedling. These are temporary and will exfoliate naturally.
- ❖ *For more details, please see the Morpheus Post Care sheet.*

### **Contact the office immediately if you have any of the following:**

- ❖ Fever of 101.5 or greater, Severe pain, blistering, changes in skin sensation

**Please contact us at (323) 909-9434 if you have any questions or concerns.**